

Amendments to the Claims:

This listing of claims will replace all prior versions, and listings, of claims in the application.

Listing of Claims:

1. – 48. (Previously Cancelled)

49. (Currently Amended) An array comprising two or more nucleic acid molecules immobilized on a substrate, wherein at least two of the nucleic acid molecules have a nucleic acid sequence consisting of:

- (a)—the nucleic acid sequence as shown in SEQ ID NOS:12, 15, 21, 22, 23, 24, 25, 26, 35 or 44;
- (b)—~~a nucleic acid sequence prepared using amplification and primer pairs, wherein the primer pairs are selected from the following pairs of nucleic acid sequences:~~

SEQ ID NO:70 and SEQ ID NO:71;
SEQ ID NO:76 and SEQ ID NO:77;
SEQ ID NO:88 and SEQ ID NO:89;
SEQ ID NO:90 and SEQ ID NO:91;
SEQ ID NO:92 and SEQ ID NO:93;
SEQ ID NO:94 and SEQ ID NO:95;
SEQ ID NO:96 and SEQ ID NO:97;
SEQ ID NO:98 and SEQ ID NO:99;
SEQ ID NO:116 and SEQ ID NO:117; or
SEQ ID NO:134 and SEQ ID NO:135;

- (c)—~~the nucleic acid sequence of (a) or (b), wherein T can be U; or~~
- (d)—~~a fragment of (a) to (c) that specifically hybridizes to one ABC transporter gene.~~

50. (Previously Amended) The array according to claim 49, wherein the array is a microarray.

51 – 72. (Previously Cancelled)

73-77. (Cancelled Herein).

78. (Currently Amended) An array for screening a sample for the presence of nucleic acid molecules that encode human ABC transporters, the array comprising a substrate having immobilized in distinct spots thereon at least 10 nucleic acid probes, wherein 10 of the probes consist of:

- 1) a probe that specifically hybridizes to a nucleic acid sequence encoding human ABC transporter B1, wherein the nucleic acid sequence of the probe is selected from the group consisting of:
 - (a) a nucleic acid sequence consisting of SEQ ID NO. 12,-
 - (b) a nucleic acid sequence prepared using amplification and primer pairs having the nucleic acid sequence of SEQ ID NO. 70 and SEQ ID NO. 71,-
 - (c) a nucleic acid sequence of a) or b) wherein T can be U, and
 - (d) a fragment of a), b) or c) that specifically hybridizes to a nucleic acid sequence encoding human ABC transporter B1;
- 2) a probe that specifically hybridizes to a nucleic acid sequence encoding human ABC transporter B4, wherein the nucleic acid sequence of the probe is selected from the group consisting of:
 - (a) a nucleic acid sequence consisting of SEQ ID NO. 15,-
 - (b) a nucleic acid sequence prepared using amplification and primer pairs having the nucleic acid sequence of SEQ ID NO. 76 and SEQ ID NO. 77,-
 - (c) a nucleic acid sequence of a) or b) wherein T can be U, and
 - (d) a fragment of a), b) or c) that specifically hybridizes to a nucleic acid sequence encoding human ABC transporter B4;
- 3) a probe that specifically hybridizes to a nucleic acid sequence encoding human ABC transporter B11, wherein the nucleic acid sequence of the probe is selected from the group consisting of:
 - (a) a nucleic acid sequence consisting of SEQ ID NO. 21,-
 - (b) a nucleic acid sequence prepared using amplification and primer pairs having the nucleic acid sequence of SEQ ID NO. 88 and SEQ ID NO. 89,-
 - (c) a nucleic acid sequence of a) or b) wherein T can be U, and
 - (d) a fragment of a), b) or c) that specifically hybridizes to a nucleic acid sequence encoding human ABC transporter B11;

- 4) a probe that specifically hybridizes to a nucleic acid sequence encoding human ABC transporter C1, wherein the nucleic acid sequence of the probe is selected from the group consisting of:
 - (a) a nucleic acid sequence consisting of SEQ ID NO. 22,-
 - (b)-a-nucleic-acid-sequence-prepared-using-amplification-and-primer-pairs having the nucleic acid sequence of SEQ ID NO. 90 and SEQ ID NO. 91,-
 - (c)-a-nucleic-acid-sequence-of-a)-or-b)-wherein-T-can-be-U,-and-
 - (d)-a-fragment-of-a),-b)-or-c)-that-specifically-hybridizes-to-a-nucleic-acid sequence-encoding-human-ABC-transporter-C1;
- 5) a probe that specifically hybridizes to a nucleic acid sequence encoding human ABC transporter C2, wherein the nucleic acid sequence of the probe is selected from the group consisting of:
 - (a) a nucleic acid sequence consisting of SEQ ID NO. 23,-
 - (b)-a-nucleic-acid-sequence-prepared-using-amplification-and-primer-pairs having the nucleic acid sequence of SEQ ID NO. 92 and SEQ ID NO. 93,-
 - (c)-a-nucleic-acid-sequence-of-a)-or-b)-wherein-T-can-be-U,-and-
 - (d)-a-fragment-of-a),-b)-or-c)-that-specifically-hybridizes-to-a-nucleic-acid sequence-encoding-human-ABC-transporter-C2;
- 6) a probe that specifically hybridizes to a nucleic acid sequence encoding human ABC transporter C3, wherein the nucleic acid sequence of the probe is selected from the group consisting of:
 - (a) a nucleic acid sequence consisting of SEQ ID NO. 24,-
 - (b)-a-nucleic-acid-sequence-prepared-using-amplification-and-primer-pairs having the nucleic acid sequence of SEQ ID NO. 94 and SEQ ID NO. 95,-
 - (c)-a-nucleic-acid-sequence-of-a)-or-b)-wherein-T-can-be-U,-and-
 - (d)-a-fragment-of-a),-b)-or-c)-that-specifically-hybridizes-to-a-nucleic-acid sequence-encoding-human-transporter-C3;
- 7) a probe that specifically hybridizes to a nucleic acid sequence encoding human ABC transporter C4, wherein the nucleic acid sequence of the probe is selected from the group consisting of:
 - (a) a nucleic acid sequence consisting of SEQ ID NO. 25,-
 - (b)-a-nucleic-acid-sequence-prepared-using-amplification-and-primer-pairs having the nucleic acid sequence of SEQ ID NO. 96 and SEQ ID NO. 97,-
 - (c)-a-nucleic-acid-sequence-of-a)-or-b)-wherein-T-can-be-U,-and-

(d) a fragment of a), b) or c) that specifically hybridizes to a nucleic acid sequence encoding human ABC transporter C4;

8) a probe that specifically hybridizes to a nucleic acid sequence encoding human ABC transporter C5, wherein the nucleotide sequence of the probe is selected from the group consisting of:-

- (a) a nucleic acid sequence consisting of SEQ ID NO. 26,-
- (b) a nucleic acid sequence prepared using amplification and primer pairs having the nucleic acid sequence of SEQ ID NO. 98 and SEQ ID NO. 99,-
- (c) a nucleic acid sequence of a) or b) wherein T can be U, and-
- (d) a fragment of a), b) or c) that specifically hybridizes to a nucleic acid sequence encoding human ABC transporter C5;

9) a probe that specifically hybridizes to a nucleic acid sequence encoding human ABC transporter D1, wherein the nucleic acid sequence of the probe is selected from the group consisting of:-

- (a) a nucleic acid sequence consisting of SEQ ID NO. 35,-
- (b) a nucleic acid sequence prepared using amplification and primer pairs having the nucleic acid sequence of SEQ ID NO. 116 and SEQ ID NO. 117,-
- (c) a nucleic acid sequence of a) or b) wherein T can be U, and-
- (d) a fragment of a), b) or c) that specifically hybridizes to a nucleic acid sequence encoding human ABC transporter D4; and

10) a probe that specifically hybridizes to a nucleic acid sequence encoding human ABC transporter G2, wherein the nucleic acid sequence of the probe is selected from the group consisting of:-

- (a) a nucleic acid sequence consisting of SEQ ID NO. 44,-
- (b) a nucleic acid sequence prepared using amplification and primer pairs having the nucleic acid sequence of SEQ ID NO. 134 and SEQ ID NO. 135,-
- (c) a nucleic acid sequence of a) or b) wherein T can be U, and-
- (d) a fragment of a), b) or c) that specifically hybridizes to a nucleic acid sequence encoding human ABC transporter G2.

79.-85. (Cancelled Herein)